

Boundary conditions and generation of periodic noise by a space-charge wave

L. P. Gor'kov

L. D. Landau Institute of Theoretical Physics, Academy of Sciences of the USSR

(Submitted 9 June 1983)

Pis'ma Zh. Eksp. Teor. Fiz. **38**, No. 2, 76–79 (25 July 1983)

The excitation of oscillations by a moving space-charge wave is attributed to a nonlinear periodic regime which arises near a contact with a normal conductor. This regime corresponds to either a homogeneous phase slippage (thin samples) or a space-charge wave which is moving across the track of edge dislocations in the lattice. Some simple microscopic equations are proposed.

PACS numbers: 72.70. + m

The excitation of nearly periodic oscillations by a space-charge wave was observed originally in NbSe₃ (Refs. 1 and 2) and later in TaS₃ (Ref. 3) and K_{0.3}MoO₃ (Ref. 4). Most of the corresponding models^{5–14} interpret the effect on the basis of a nonlinear equation for the phase of the space-charge wave in the presence of a pinning potential, in an effort to relate the excitation effect to the presence of a threshold electric field (E_t) required for motion of the wave. In the present letter we argue that when the pinning of the space-charge wave is a volume effect (defects¹⁵) the excitation of periodic oscillations results from the boundary conditions near a contact with a normal conductor, which give rise to phase-slippage centers. This mechanism, from the theory of superconductivity,^{16,17} requires incorporating the change in the modulus of the order parameter.

Space-charge waves are described by a lattice strain $u(x) = u_0 \cos(2k_F x + \phi)$, where an arbitrary phase in the incommensurate case reflects the possibility that the wave can move in the absence of pinning. We will ignore the latter ($E > E_t$). As the order parameter

$$\Delta(x, t) = |\Delta(x, t)| \exp i\phi(x, t) \quad (1)$$

in the slowly moving fields we adopt the size of the “gap” in the electron spectrum, which is proportional to the local strain.

A volume strain (1) can occur only below the transition temperature T_p , while a large strain (large on the atomic scale) is unavoidable near an interface between two media. This strain is relatively insensitive to the low-temperature structural transition. If it were not, a structure appearing spontaneously in the volume at $T < T_p$ would have to change over a distance $\xi_0 \sim \hbar v / T_p$ to suit the existing strain at the boundary. This condition unavoidably fixes the phase of the space-charge wave at the boundary.

Assuming that the change $|\Delta|$ is small in a weak field, we find

$$\Gamma\phi = aE + \xi_0^2 \nabla^2 \phi. \quad (2)$$

Ordinarily (the relaxation time τ is set by impurities) we would have^{18,19}

$$a \sim e \xi_0 / T_p, \quad \Gamma^{-1} \sim T_p^2 \tau. \quad (3)$$

A steady-state ($\dot{\phi} = 0$) solution of (2) can occur in the volume only in the presence of pinning forces; in the opposite case, vector (1) rotates at a frequency

$$\Omega = a \Gamma^{-1} E \sim eEl, \quad (4)$$

corresponding to the motion of a space-charge wave at a velocity $2k_F u = \Omega$.

Figure 1 is a mechanical model described by (2). A spring fixed at $x = 0$ is rotated at a distant point at the angular velocity Ω given by (4). A steady-state process can be established if, in some cross section $x = x_0$, the coupling between two regions is periodically broken because of the accumulated stress, and a phase slippage $[\phi] = 2\pi$ occurs. In parabolic equation (2) the term with the field is a spatially homogeneous source, while the slippage process provides a phase "sink": $2\pi\Gamma^{-1}\delta(x - x_0)\delta(t - t_n)$, where $t_n = 2\pi\Omega^{-1}n$.

The arguments can be supported by a convenient microscopic model. We assume that the electron spectrum $\epsilon(\mathbf{p})$ is three-dimensional and corresponds to two regions with the property of "superposition,"²⁰ $\epsilon(\mathbf{p} + \mathbf{q}) = -\epsilon(\mathbf{p})$ (for example, tunneling between chains corresponds to the strong-coupling approximation²¹). If the longitudinal component $q_{\parallel} = 2k_F$ is incommensurate, the metal-dielectric transition gives rise to superstructure (1). Particularly simple equations arise if the transition temperature is lowered ($T_p \ll T_{p0}$) by defects, i.e., in the range of concentrations below the critical value $1/\tau_c = 4\gamma/3\pi T_{p0}$:

$$\Delta + (\Delta^2 + Q^2 - 1)\Delta - \nabla^2 \Delta = 0, \quad (5)$$

$$(E + \phi)\Delta^2 - \text{div}(Q\Delta^2) = 0$$

(where $\Delta \equiv |\Delta|$, $Q = \nabla\phi$). The order parameter in (5) is expressed in units of $\Delta_{\infty} = \pi T_p \delta^{1/2}$ ($\delta = 1 - T^2/T_p^2$); the field eE is expressed in units of $\pi^2 T_p^2 \delta / 12\bar{v}$; and the times and coordinates are expressed in units of $9/2\pi^2 T_p^2 \tau_c \delta$ and $\sqrt{3\bar{v}^2/2}/\pi T_p$, respectively. The current is carried primarily by normal carriers,

$$j = \frac{\sigma_n \pi^2 T_p^2}{12 e \bar{v}} \left\{ E - \lambda E \Delta^2 - \left(\frac{8}{9} \right) \lambda \epsilon \Delta^2 \dot{\phi} \right\}. \quad (6)$$




FIG. 1.

The second term in (6) is a small correction to the normal conductivity caused by the transition, while the third term reflects the moving space-charge wave ($\lambda = (8\pi^2/3)T_p^2\tau_c^2\delta$, $\epsilon = \bar{v}^2/\bar{v}^2$).

The point x_0 in Fig. 1 is determined by integrating nonlinear equations (5) and using the conditions $\Delta = \Delta_0$ and $\phi = 0$ at $x = 0$ and $\Delta = 1$ and $\dot{\phi} = -E$ in the limit $x \rightarrow \infty$. The choice of x_0 is not obvious. At $E \ll 1$ we could have a regime with $x_0 \gg 1$. At $x < x_0$, all quantities would then be periodic in the time, and Δ would vary slowly along the coordinate. For the average quantities we find from (5)

$$\bar{\Delta}^2 = 1 - \bar{Q}^2; \quad 3\bar{Q} - 2 \operatorname{Arctg} \bar{Q} = Ex.$$

Corresponding to the stable branch of the solutions of the second equation is the value $\bar{Q} < 1/3$:

$$Ex_0 < 1 - \ln 2. \quad (7)$$

This example shows that an electric field creates near the boundary a phase gradient, i.e., a local change in the period of the space-charge wave. This change increases along the coordinate (and over time) until the superstructure loses its stability. At distances $|x - x_0| \sim 1$, rapid changes occur in all quantities, and Δ crosses zero.

At $E \gg 1$ the solution yields $x_0 \cong (2E)^{-1/2} \ln E$. The amplitude of the variable stress ($j = \text{const}$) is

$$e V_1 = \left(1 - \frac{8}{9}\epsilon\right) \frac{2\pi^3}{3\sqrt{3}} \epsilon^{1/2} \delta(T_p\tau_c)^2 T_p \sin(24 eEt).$$

In a sample which is unbounded in the transverse direction a phase sink can occur in the x_0 plane because of the motion of a periodic structure of the phase-slippage centers if the period L and velocity v_1 of the structure are related by $2\pi v_1 = \Omega L$. An estimate from (5) would give us $v_1 \sim v(T_p\tau_c)\delta$. Physically, this structure would represent edge dislocations in the lattice of the space-charge wave. Which solution to choose (that corresponding to the homogeneous phase-slippage center or that corresponding to a traveling track) is not clear. Fung and Steeds²³ have shown that an NbSe₃ single crystal consists of normal domains ($\sim 10^{-6}$ cm wide) oriented parallel to the principal axis; this situation would seem to further hinder the motion of dislocations.

According to (4), the frequency of the oscillations which are excited is a measure of the mean free path in actual substances. In turn, working from (2) and (3) we find the following criterion according to result (7) (under the condition $\xi_0 Q \sim 1$):

$$e E x_0 \sim T_p.$$

We then find $x_0 \sim 10^{-2}$ cm (with $E \sim 100$ mV/cm and $T_p \sim 50$ K). In other words, x_0 is approximately equal to the length $l \sim 100 \mu\text{m}$ found in Ref. 22 from an estimate of the effect of the sample geometry.

It should be noted that the concept of phase-slippage centers was recently proposed independently in an experimental study by Ong and Verma²⁴ in an effort to explain data on the local nature of the excitation mechanism in NbSe₃.

I wish to thank S. A. Brazovskii and N. B. Kopnin for useful discussions and B. I. Ivlev for useful comments.

- ¹R. M. Fleming and C. G. Grimes, *Phys. Rev. Lett.* **42**, 1423 (1979).
- ²P. Monceau, J. Richard, and M. Renard, *Phys. Rev. Lett.* **45**, 43 (1980); *Phys. Rev. B* **25**, 931 (1982).
- ³G. Grüner, A. Zettl, W. G. Clark, and A. H. Thompson, *Phys. Rev. B* **23**, 6813 (1981).
- ⁴J. Dumas, C. Schlenker, J. Marcus, and R. Buder, *Phys. Rev. Lett.* **50**, 757 (1983).
- ⁵J. Bardeen, *Phys. Rev. Lett.* **45**, 1978 (1980).
- ⁶M. Weger, G. Grüner, and W. G. Clark, *Solid State Commun.* **35**, 243 (1980).
- ⁷G. Grüner, A. Zawadowski, and P. M. Chaikin, *Phys. Rev. Lett.* **46**, 511 (1981).
- ⁸S. E. Burkov, V. L. Pokrovsky, and G. V. Uimin, *Solid State Commun.* **40**, 363 (1981).
- ⁹P. Bak, *Phys. Rev. Lett.* **48**, 692 (1982).
- ¹⁰S. N. Artemenko and A. F. Volkov, *Pis'ma Zh. Eksp. Teor. Fiz.* **33**, 155 (1981) [*JETP Lett.* **33**, 147 (1981)].
- ¹¹M. Weger and B. Horovitz, *Solid State Commun.* **43**, 583 (1982).
- ¹²L. Sneddon, M. C. Cross, and D. S. Fisher, *Phys. Rev. Lett.* **49**, 292 (1982).
- ¹³R. M. Klemm and J. R. Schrieffer, Preprint, 1983.
- ¹⁴S. E. Banes and Z. Zawadowski, Preprint, 1983.
- ¹⁵P. A. Lee and T. M. Rice, *Phys. Rev. B* **19**, 3970 (1979).
- ¹⁶J. S. Langer and V. Ambegaokar, *Phys. Rev.* **164**, 498 (1967).
- ¹⁷B. I. Ivlev and N. B. Kopnin, *Pis'ma Zh. Eksp. Teor. Fiz.* **28**, 640 (1978) [*JETP Lett.* **28**, 592 (1978)].
- ¹⁸L. P. Gor'kov and E. N. Dolgov, *Zh. Eksp. Teor. Fiz.* **77**, 396 (1979) [*Sov. Phys. JETP* **50**, 203 (1979)]; *J. Low Temp. Phys.* **24**, 101 (1981).
- ¹⁹S. N. Artemenko and A. F. Volkov, *Zh. Eksp. Teor. Fiz.* **81**, 1872 (1981) [*Sov. Phys. JETP* **54**, 992 (1981)].
- ²⁰L. V. Keldysh and Yu. V. Kopaev, *Fiz. Tverd. Tela (Leningrad)* **6**, 2791 (1964).
- ²¹B. Horovitz, H. Gutfreund, and M. Weger, *Solid State Commun.* **39**, 541 (1981).
- ²²J. C. Gill, *Solid State Commun.* **44**, 1041 (1982).
- ²³K. K. Fung and J. W. Steeds, *Phys. Rev. Lett.* **45**, 1696 (1980).
- ²⁴N. P. Ong and G. Verma, *Phys. Rev. B* **27**, 4495 (1983); N. P. Ong, G. Verma, and K. Maki, Preprint, 1983.

Translated by Dave Parsons

Edited by S. J. Amoretty